



Clinical epidemiological and hematological characterization of hospitalized patients with clinical suspicion of dengue

Caracterización clínico epidemiológica y hematológica de pacientes hospitalizados con sospecha clínica de dengue

Yoánder Pérez Díaz¹ <https://orcid.org/0000-0003-3439-7424>

Rolando Rodríguez Puga^{1*} <https://orcid.org/0000-0003-3350-374X>

Melissa María Navarro Gómez² <https://orcid.org/0009-0009-0053-7338>

Yanira González Ronquillo² <https://orcid.org/0000-0002-2177-4297>

Luis Acao Francois² <https://orcid.org/0000-0001-5037-1027>

Yanet Pérez Guerrero³ <https://orcid.org/0000-0002-1306-7154>

*Corresponding author (email): rolandote1986@gmail.com

¹University of Medical Sciences of Camagüey. School of Medicine. Provincial Pediatric Teaching Hospital “Dr. “Eduardo Agramonte Piña.” Department of Hospital Epidemiology. Camagüey, Cuba.

²University of Medical Sciences of Camagüey. School of Medicine. Department of Public Health. Camagüey, Cuba.

Abstract

Introduction: Dengue is one of the main viral diseases of epidemic nature. The study of the clinical, epidemiological and hematological characteristics is essential for the prevention, treatment and evolution of cases.

Objective: To characterize hospitalized patients with clinical suspicion of dengue from a clinical, epidemiological and hematological point of view.

Methods: A descriptive, cross-sectional study was carried out at the “Amalia Simoni” Provincial Hospital of Camaguey, Cuba, during the year 2022. The sample was made up of 157 hospitalized patients with clinical suspicion of dengue. The variables analyzed were: age group, sex, month of diagnosis, signs and symptoms, hematological variables, leukocyte differential formula and days of onset of symptoms.

Results: There was superiority in the age group of 40-49 years (28.7 %) and in the male sex (50.3 %). In the month of August, 33 patients (21.0 %) were reported, while the main signs and symptoms were fever (90.4 %) and headache (59.9 %). Regarding hematological variables, the mean hematological variables,

the mean hematocrit ranged between 0.42 and 0.48 L/l and the platelet count between 224 and 256 x 10⁹/L. In relation to the differential formula, the average number of polymorphonuclear cells varied between 0.62 and 0.72 % and that of lymphocytes between 0.28 and 0.36 %.

Conclusions: The sample studied was characterized by the preponderance of patients in the fourth decade of life who presented hyperthermia as a fundamental clinical manifestation. The study of hematological variables was essential for the evolution of the cases.

Keywords: dengue; virus; prevention; platelets; hematocrit.

Resumen

Introducción: El dengue es una de las principales enfermedades virales de naturaleza epidémica. El estudio de las características clínicas, epidemiológicas y hematológicas es fundamental para la prevención, tratamiento y evolución de los casos.

Objetivo: Caracterizar desde el punto de vista clínico epidemiológico y hematológico a pacientes hospitalizados con sospecha clínica de dengue.

Métodos: Se realizó un estudio descriptivo, de corte transversal, en el Hospital Provincial Clínico Quirúrgico Docente “Amalia Simoni” de Camagüey, durante el año 2022. La muestra quedó conformada por 157 pacientes hospitalizados con sospecha clínica de dengue. Las variables analizadas fueron: grupo de edad, sexo, mes del diagnóstico, signos y síntomas, variables hematológicas, fórmula diferencial de leucocitos y días de inicio de los síntomas.

Resultados: Existió superioridad del grupo de edad de 40-49 años (28,7 %) y del sexo masculino (50,3 %). En el mes de agosto se reportaron 33 pacientes (21,0 %), mientras los principales signos y síntomas fueron la fiebre (90,4 %) y la cefalea (59,9 %). En cuanto a las variables hematológicas la media del hematocrito osciló entre 0,42 y 0,48 L/l y el conteo de plaquetas entre 224 y 256 x 10⁹/L. En relación con la fórmula diferencial el promedio de polimorfonucleares varió entre 0,62 y 0,72 % y el de linfocitos entre 0,28 y 0,36.

Conclusiones: La muestra estudiada se caracterizó por la preponderancia de pacientes en la cuarta década de vida que presentaron hipertermia como manifestación clínica fundamental. El estudio de las variables hematológicas resultó indispensable para la evolución de los casos.

Palabras clave: Dengue; virus; prevención; plaquetas; hematocrito.

Introduction [1]

Dengue is a viral infection transmitted by mosquitoes, highly prevalent in tropical and subtropical regions, and responsible for putting more than four billion individuals at risk globally. It is estimated that about 500,000 people with the severe form of dengue require hospitalization every year, including the pediatric group. 2.5% of the hospitalized patients dies from complications associated with the severe form of the disease.^(1,2)

Dengue is an arbovirus that has constantly expanded to new areas and territories where control had been achieved. Currently, dengue has spread to more than 129 countries, and cases are increasing steadily

over time. About 70 % of the global burden of the disease is concentrated in Asia. In particular, countries in the Southeast Asian Region are highly affected, with an estimated of 52 % of the world's population at risk of becoming ill.^(2,3)

About half of the planet's inhabitants currently live in environments conducive to the transmission of dengue. Infection occurs through mosquitoes of the *Aedes* species, which proliferate in tropical and subtropical areas. Combined with these global trends, rising temperatures attributed to climate change have raised concerns that dengue will intensify in already endemic areas through more rapid viral amplification.^(4,5)

In the Region of the Americas, it is the arbovirus that causes the greatest number of cases, with epidemics that occur cyclically every 3 to 5 years. In 2019, the highest historical number of notifications was recorded, with more than 3.1 million cases, of which 28,203 were serious and 1,773 deaths. During the year 2022, until epidemiological week (EW) 52, of the 2,803,096 cases reported in the region, 1,299,273 (46.4 %), were confirmed by laboratory tests.^(3,6)

Since 2020, the circulation of dengue, Chikungunya and Zika viruses occurs simultaneously with the active transmission of the SARS-CoV-2 virus in endemic countries and territories, in the Region of the Americas. Health systems must remain alert to the coexistence of these two clinical entities due to the population effect that it may cause.⁽⁷⁾

The identification of new SARS-CoV-2 variants of concern and inadequate coverage of the COVID-19 vaccine, in endemic areas of these arboviruses where public health and social measures to prevent COVID-19 have been made more flexible, pose a complex epidemiological situation, a high demand on health services, as well as a constant challenge for health systems in all their components and levels, including diagnosis, epidemiological surveillance and management of these cases.^(8,9)

The region of the Americas is today a geographical area with a high impact of arboviruses, which is why the countries of the region have strengthened their health systems for prevention and control, with the establishment of comprehensive surveillance systems (clinical surveillance epidemiological, laboratory, entomological and environmental). Cuba does not escape the risk, which is why confronting dengue and other arboviruses is a priority for the country's highest authorities.⁽⁸⁾

Cuba throughout history has shown high levels of *Aedes aegypti* infestation, so that, between 1977 and 1978, it reported the first dengue epidemic, corresponding to the second half of the 20th century, and caused by the dengue virus (DENV-1). This was characterized as an epidemic of dengue fever (DF), with more than 400 thousand cases from all over the country. Seroepidemiological studies showed that 44.46 % of the population was infected by this virus.⁽⁹⁾

In 1981, Cuba reported the first epidemic of Dengue Hemorrhagic Fever (DH) in the region, with more than 300 thousand cases, 10 thousand serious and 158 deaths, of them, 101 children. The epidemic,

caused by DENV-2, affected the entire country. The transmission was stopped after an intense vector control campaign that lasted 4 months, until its elimination was achieved. The country was free of transmission from 1982 to 1996. In 1997, the Santiago de Cuba municipality reported an epidemic of DHF caused by DENV-2, with more than 5 thousand cases.⁽¹⁰⁾

In the years 2001-2002, a new DH epidemic occurred, caused by DENV-3, which greatly affected the City of Havana. This epidemic, which began in June 2001, was preceded by a small outbreak due to DENV-3 and 4 in health areas of the Boyeros, Playa and Lisa municipalities at the end of 2000.⁽⁸⁾

In 2005, three small outbreaks of DENV-3 and 4 occurred in Havana City and Camagüey, eliminated in a timely manner. In 2006, an epidemic occurred, controlled at the beginning of 2007, while in the period 2007-2022, several outbreaks were recorded throughout the country, with circulation of several viral serotypes.^(9,10) For the reasons described above, the present research with the objective of characterizing hospitalized patients with clinical suspicion of dengue from a clinical, epidemiological and hematological point of view.

Methods [1]

A descriptive, cross-sectional investigation was carried out at the “Amalia Simoni” Provincial Hospital in Camagüey, Cuba, during the year 2022. The sample was made up of 157 hospitalized patients with clinical suspicion of dengue, selected using the eligibility criteria.

Inclusion criteria

- Patient with a presumptive diagnosis of dengue at the time of admission and a definitive diagnosis during hospital discharge.
- Patient aged 19 years or older.

Exclusion criteria

- Patient rediscussed due to presenting a diagnosis other than the clinical suspicion of dengue raised during hospitalization.
- Patient with incomplete medical history, so that it was impossible to consult the values of the variables at the different cut-off points evaluated in the study.

The variables were the following:

- Age group (19-29, 30-39, 40-49, 50-59, 60-69 and ≥ 70).
- Sex (male and female).
- Month of diagnosis (January, February, March, April, May, June, July, August, September, October, November and December).
- Hematological variables (hematocrit, hemoglobin, platelets and leukocytes).
- Day of hospital stay (1,2,3,4,5,6 and 7).
- Differential formula (polymorphonuclear cells, lymphocytes, eosinophils and monocytes).

The reference values for hematocrit in men are between 0.41-0.54 L/l and in women: 0.37-0.47 L/l. The normal global leukocyte count should range between 5-10 x 10⁹/L, the platelet counts between 150-450 x 10⁹/L and polymorphonuclear cells between 0.55-0.65%. Similarly, the allowable numbers of lymphocytes are between 0.25-0.35%, monocytes between 0.03-0.06 % and eosinophils between 0.02-0.04%.

The statistical registry and the patients' medical records were the source of information. The data were entered into an Excel database for processing, obtaining absolute numbers and percentages. The results were presented in the form of texts and a comparison was made with those achieved by national and international authors, which allowed conclusions to be reached. Descriptive statistics were applied to obtain absolute numbers and percentages.

The study was approved by the Ethics Committee and the Scientific Council of the “Amalia Simoni” Provincial Hospital of Camagüey. The confidentiality of the data was maintained by coding the variables, accessible only to the researchers and used for scientific purposes. The bioethical principles contained in the Helsinki Declaration were taken into account.⁽¹¹⁾

Results [1]

Table 1 shows the group of 40-49 years of age as the group with the highest prevalence of cases (28.7%), while the group with the lowest incidence was those ≥ 70 years of age. The male sex resulted in a discreet preponderance in relation to the female sex (79/78).

Table 1 - Distribution of patients with dengue according to age group and sex

Age group (years)	Sex				Total		Measures	
	Female		Male					
	No.	%	No.	%	No.	%	Half	D.S
19-29	17	10,8	18	11,5	35	22,3	23,4	5,12
30-39	5	3,2	10	6,4	15	9,6	34,2	4,98
40-49	27	17,2	18	11,5	45	28,7	43,6	5,22
50-59	18	11,5	17	10,8	35	22,3	56,1	5,34
60-69	9	5,7	10	6,4	19	12,1	64,7	4,78
≥70	2	1,3	6	3,7	8	5,0	76,6	5,67
Total	78	49,7	79	50,3	157	100,0	49,8	6,03

Source: Patient's clinical chart

Regarding the total number of cases by month, figure 1 shows an increase in the month of March (24), with a decrease in the months of April and May, and then experiencing a sharp drop in the month of

June (3). The highest notification of cases occurred in the month of August (33), while the remaining four months had similar behavior.




Fig. 1 – Distribution of patients with dengue according to month of diagnosis.
Source: Patient’s clinical chart

The main signs and symptoms (Fig. 2) were represented by fever in 90.4% of cases, followed by headache (59.9%) and arthralgia (50.3%). Abdominal pain was the most prominent warning sign (16.6%).




Fig. 2 – Distribution of dengue patients according to signs and symptoms.
Source: Patient’s clinical chart

Note: In Fig. 2, the percentage was calculated in relation to the total sample (157), so that the sum of the clinical manifestations does not reach 100%, since each patient presented more than one sign or symptom.

Table 2 analyzes the average of the main hematological variables. Hematocrit reached the lowest values during the first day of hospital stay and the highest values on the seventh day. Hemoglobin remained with minimal variations during the course of the seven days of monitoring, while the lowest average platelet count occurred between the third and fourth day, with recovery on the seventh day. Leukopenia became noticeable between the fifth and sixth day.

Table 2 - Average values of hematological variables according to days of hospital stay in patients with dengue

Hematological variables	Day of hospital stay						
	1	2	3	4	5	6	7
	Average						
Hematocrit L/l	0,42	0,43	0,43	0,43	0,43	0,43	0,48
Hemoglobin g/L	13,4	13,5	13,4	13,5	13,6	13,4	13,5
Platelets x 10 ⁹ /L	240	231	225	224	230	240	256
Leukocytes x 10 ⁹ /L	8,6	7,4	6,2	6,3	5,7	5,8	6,1

Source: Patient's clinical chart

In the differential formula (Table 3), during the first day of hospital stay the polymorphonuclear cells reached the maximum percentage (0.72), reaching normal numbers on the seventh day. Lymphocytes were lower during the first day and then remained little variable, eosinophils showed no changes, while monocytes increased on the sixth day.

Table 3 - Average values of the leukocyte differential formula according to days of hospital stay in patients with dengue

Differential formula	Day of hospital stay						
	1	2	3	4	5	6	7
	Average						
PMN	0.72	0.66	0.67	0.62	0.64	0.62	0.62
Lymphocytes	0.28	0.33	0.34	0.36	0.36	0.36	0.35
Eosinophils	0.00	0.00	0.01	0.01	0.01	0.01	0.00
Monocytes	0.01	0.03	0.03	0.04	0.04	0.05	0.02

Source: Patient's clinical chart

PMN: Polymorphonuclear

Discussion [1]

The epidemiological history and clinical manifestations are of vital importance to establish the presumptive diagnosis of dengue. Hematological examinations complement the diagnostic orientation and allow the evolution of cases. For this reason, the present study is developed in which a series of variables are characterized that will be the subject of discussion.

In relation to age, similar results were found to those obtained by authors such as Adane and Getawa,⁽¹²⁾ Araya and Wilson,⁽¹³⁾ and Khan et al.⁽¹⁴⁾ In this same aspect, Beltrán Saiz et al,⁽¹⁹⁾ They identified the group of 41-50 years as the most representative, while the predominant sex was male with very discrete figures. Similar results were found in the present investigation.

Other authors,^(15,16) also found a higher prevalence of infection in men. It is the opinion of those responsible for this research that it may be due to cultural habits, where men often take off their shirts during the hot summer months and spend the entire night exposed to mosquito bites.

The month with the highest frequency of cases was August, a result that agrees with that obtained by Khan et al.⁽¹⁴⁾ in Pakistan. Garcia et al.⁽¹⁵⁾ assert that the risk of dengue infection is influenced by rainfall, temperature and the degree of urbanization.

In the opinion of the authors of the study, the highest incidence of dengue in the month of August could be related to the temperatures and humidity that favor the reproduction of the vector, in addition to abundant rainfall. Due to frequent power outages in the summer months due to high demand, people store water in various containers that serve as larval breeding habitats, resulting in an increase in mosquito population density.

Fever is a clinical manifestation that accompanies a large number of diseases, highlighting infectious diseases, which particularly include dengue. Khan et al.⁽¹⁴⁾ conducted a community-based cross-sectional study where they reported fever and bone pain as fundamental symptoms, followed by headache; Similar results were found in the present work.

In a systematic review,⁽⁵⁾ on the prevalence and distribution of dengue, reference was made to the high seroprevalence in correspondence with the report of fever, results with which the present investigation coincides. Luque et al.⁽¹⁶⁾ observed in their series of cases that the most frequent clinical manifestations were related to the febrile phase of dengue: fever and systemic symptoms such as headache and nausea.

Hematological alterations are common in the course of dengue infection, given the pathophysiology of this disease, most frequently found to be a decrease in the number of platelets (thrombocytopenia) and leukocytes (leukopenia), as well as an increase in hematocrit. The appearance of these laboratory

findings is frequently reported in various series and usually occur between the third and eighth day of the course of the disease.⁽¹⁷⁾

When evaluating the evolution and behavior of the values of the hematological variables in the study series in question, it was possible to verify variations that are related to those reported in the literature. Hematocrit is a sign of hemoconcentration secondary to hypovolemia and a variation equal to or greater than 20 % with respect to an initial value or a decrease during convalescence with respect to the critical period is considered diagnostic of dengue hemorrhagic fever.⁽¹⁸⁾

In the literature, a decrease in the number of leukocytes is mentioned as the first abnormality in the blood count. The overall decrease in leukocyte count observed in dengue patients is primarily due to a decrease in the granulocyte population (e.g. neutrophils). The ability of the dengue virus to suppress the production of white blood cells in the bone marrow may explain the appearance of leukopenia.⁽¹⁹⁾

Leukopenia is usually a common finding in around 68.4% of patients. The differential formula can show neutropenia, especially in the initial phase of the disease. The decrease in total leukocytes begins in the first three days and is slight and, five days after the illness, it is present in the majority of patients; In severe forms of the disease it is intense and marked. The most notable and characteristic thing in these patients is the differential picture with lymphocytosis and monocytosis.⁽²⁰⁾

When the cell is infested with the dengue virus, the infection migrates to the lymph nodes, monocytes and macrophages are recruited, which become the target of the infection, and then amplify and spread through the lymphatic system. As a result of this primary viremia, several cells of the mononuclear lineage, monocytes, myelocytes, and splenic and hepatic macrophages are affected. It is even proposed that the virus has the ability to affect mononuclear cells circulating in the blood.⁽⁹⁾

Platelets are an important blood component involved in clotting. Patients infected with dengue virus tend to develop thrombocytopenia during the course of infection, which makes them vulnerable to hemorrhagic manifestations and other serious complications.⁽²¹⁾

The authors of the current study estimate that platelet counts begin to fall during the febrile stage and reach their lowest point during the toxic stage. The development of thrombocytopenia in dengue infection could be due to bone marrow depression observed in the acute stage of dengue virus infection. Other explanations are direct infection of megakaryocytes leading to further destruction or the presence of antibodies directed against platelets. Another mechanism is the increase in platelet consumption through interaction with infected endothelial cells. An in vitro study demonstrated that some endothelial cells damaged by dengue can promote platelet adhesion and lysis.⁽²¹⁾

Currently, few studies are carried out that include the analysis of hematological aspects in patients with dengue. Its importance is essential for the evolution and monitoring of cases. The main limitation was

the impossibility of establishing causal relationships between the dengue event and the variables analyzed, typical of this type of study.

Conclusions [1]

The sample studied was characterized by the preponderance of patients in the fourth decade of life who presented hyperthermia as a fundamental clinical manifestation. The study of hematological variables was essential for the evolution of the cases.

From a medical perspective, it is recommended to avoid the administration of non-steroidal anti-inflammatory drugs, aspirin or other antiplatelet agents. The patient must stay under a mosquito net and use insect repellents to prevent mosquitoes from becoming infected and infecting other people.

References [1]

1. Sirisena PDN, Mahilkar S, Sharma C, Jain J, Sunil S. Concurrent dengue infections: Epidemiology & clinical implications. *Indian J Med Res.* 2021;154(5):669-779. Disponible en: http://dx.doi.org/10.4103/ijmr.IJMR_1219_18
2. Khan J, Ghaffar A, Khan SA. The changing epidemiological pattern of Dengue in Swat, Khyber Pakhtunkhwa. *PLoS One.* 2018;13(4):e0195706. Disponible en: <http://dx.doi.org/10.1371/journal.pone.0195706>
3. Messina JP, Brady OJ, Golding N, Kraemer MUG, Wint GRW, Ray SE, et al. The current and future global distribution and population at risk of dengue. *Nat Microbiol.* 2020;4(9):1508-15. Disponible en: <http://dx.doi.org/10.1038/s41564-019-0476-8>
4. Guzmán Tirado MG, Vázquez S, Álvarez M, Pelegrino JL, Ruiz Amores D, Martínez PA, et al. Vigilancia de laboratorio de dengue y otros arbovirus en Cuba, 1970-2017. *Rev Cubana Med Trop* [Internet]. 2020 [acceso 26/12/2023];71(1): e338. Disponible en: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0375-07602019000100008&lng=es
5. Prattay KMR, Sarkar MR, Shafiullah AZM, Islam MS, Raihan SZ, Sharmin N. A retrospective study on the socio-demographic factors and clinical parameters of dengue disease and their effects on the clinical course and recovery of the patients in a tertiary care hospital of Bangladesh. *PLoS Negl Trop Dis.* 2022;16(4):e0010297. Disponible en: <http://dx.doi.org/10.1371/journal.pntd.0010297>
6. Bhatt P, Sabeena SP, Varma M, Arunkumar G. Current Understanding of the Pathogenesis of Dengue Virus Infection. *Curr Microbiol.* 2021;78(1):17-32. Disponible en: <http://dx.doi.org/10.1007/s00284-020-02284-w>
7. Beltrán Saínz R, Hernández Baguer R, Mena Grandales K, Savón George L. Complicaciones oculares asociadas a la fiebre por dengue. *Revista Cubana de Oftalmología* [Internet]. 2020 [acceso 26/12/2023];33(2):810. Disponible en: <http://revoftalmologia.sld.cu/index.php/oftalmologia/article/view/810>.
8. Cardl Artal F. Complicaciones neurológicas asociadas a la infección por el virus del dengue. *Rev neurol* [Internet]. 2019 [acceso 26/12/2023];69:113-22. Disponible en: <https://temas.sld.cu/dengue/2019/12/01/complicaciones-neurolgicas-asociadas-a-la-infeccion-por-el-virus-del-dengue/>.

9. Rodríguez Puga R, Pérez Díaz Y, Dueñas Rodríguez Y, Rodríguez Abalo OdC, Díaz Pérez L, Pérez Díaz OA. Características dermatológico-epidemiológicas de las manifestaciones cutáneas por dengue en niños. *Folia Dermatológica Cubana* [Internet]. 2023 [acceso 26/12/2023];16(1):[aprox. 17 p.]. Disponible en: <https://revfdc.sld.cu/index.php/fdc/article/view/276>
10. Dueñas Rodríguez Y, Rodríguez Puga R. Características de pacientes con prescripción médica de Tratho Dengue. *Rev. Cub. Tecnol. Salud* [Internet]. 2023 [acceso 26/12/2023];14(1):58-67. Disponible en: <https://revtecnologia.sld.cu/index.php/tec/article/view/4037>
11. World Medical Association. Declaration of Helsinki. Ethical Principles for Medical Research Involving Human Subjects. *Clin Rev Educ* [Internet]. 2013 [acceso 26/12/2023];310(20):2191-4. Disponible en: <https://www.wma.net/wp-content/uploads/2016/11/DoH-Oct2013-JAMA.pdf>
12. Adane T, Getawa S. Coagulation abnormalities in Dengue fever infection: A systematic review and meta-analysis. *PLoS Negl Trop Dis*. 2021;18;15(8):e0009666. DOI: <http://dx.doi.org/10.1371/journal.pntd.0009666>
13. Araya Umaña L, Wilson Salazar E. Dengue hallazgos hematológicos y de imagen. *Revista Médica de Costa Rica y Centroamerica* [Internet]. 2016 [acceso 26/12/2023];12(2):[aprox. 12 p.]. Disponible en: <https://www.medigraphic.com/pdfs/revmedcoscen/rmc-2016/rmc163at.pdf>
14. Khan J, Adil M, Wang G, Tsheten T, Zhang D, Pan W, et al. A cross-sectional study to assess the epidemiological situation and associated risk factors of dengue fever; knowledge, attitudes, and practices about dengue prevention in Khyber Pakhtunkhwa Province, Pakistan. *Front Public Health*. 2022;29(10):923:37. DOI: <https://www.doi:10.3389/fpubh.2022.923277>
15. García Gómez, Rodríguez Morales O, Fernández González CT, Rodríguez Roque MO. Evolución clínico-epidemiológica de pacientes con diagnóstico de dengue. *Medisur* [Internet]. 2020 [acceso 26/12/2023];17(6):790-6. Disponible en: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S1727-897X2019000600790&lng=es
16. Luque N, Cilloniz C, Pons MJ, Donaires F, Albornoz R, Mendocilla Risco M, et al. Características clínicas y epidemiológicas de las muertes por dengue durante un brote en el norte del Perú. *Rev. perú. med. exp. salud pública*. 2023;40(1):67-72. DOI: <http://dx.doi.org/10.17843/rpmesp.2023.401.12148>
17. Anker M, Arima Y. Male-female differences in the number of reported incident dengue fever cases in six Asian countries. *Western Pac Surveill Response J*. 2011;2(2):17-23. DOI: <http://dx.doi.org/10.5365/WPSAR.2011.2.1.002>
18. González Perrota N, Araya S, Lovera D, Martínez de Cuellar C, Arbo Sosa A. Características clínicas y de laboratorio en pacientes pediátricos ambulatorios con Fiebre Dengue. *Pediatr. (Asunción)* [Internet]. 2020 [acceso 26/12/2023]; 47(1):5-10. Disponible en: http://scielo.iics.una.py/scielo.php?script=sci_arttext&pid=S1683-98032020000100005&lng=en
19. Shrestha DB, Budhathoki P, Gurung B, Subedi S, Aryal S, Basukala A, et al. Epidemiology of dengue in SAARC territory: a systematic review and meta-analysis. *Parasit Vectors*. 2022;15(1):389-96. Disponible en: <https://www.doi:10.1186/s13071-022-05409-1>
20. Chaloeuwong J, Tantiworawit A. Useful clinical features and hematological parameters for the diagnosis of dengue infection in patients with acute febrile illness: a retrospective study. *BMC Hematology* [Internet]. 2020 [acceso 26/12/2023];15(3):380-9. Disponible en:

<https://pubmed.ncbi.nlm.nih.gov/30181881/>

21. Castilho BM, Silva MT, Freitas ARR, Fulone I, Lopes LC. Factors associated with thrombocytopenia in patients with dengue fever: a retrospective cohort study. *BMJ Open*. 2020;10(9):e035120. DOI: <http://dx.doi.org/10.1136/bmjopen-2019-035120>

Dislosure

The authors declare that does not exist an interest conflict.

Author contributions

Conceptualization: Yoánderosn Pérez Díaz, Rolando Rodríguez Puga and Melissa María Navarro Gómez.

Data curation: Yoánderosn Pérez Díaz, Rolando Rodríguez Puga, Melissa María Navarro Gómez and Yanet Pérez Guerrero.

Formal analysis: Yoánderosn Pérez Díaz and Rolando Rodríguez Puga.

Research: Yoánderosn Pérez Díaz and Rolando Rodríguez Puga.

Methodology: Yoánderosn Pérez Díaz and Rolando Rodríguez Puga.

Project administration: Yoánderosn Pérez Díaz and Rolando Rodríguez Puga.

Supervision: Yoánderosn Pérez Díaz, Luis Acao Francois and Yanira González Ronquillo.

Validation: Yoánderosn Pérez Díaz, Luis Acao Francois and Yanira González Ronquillo.

Visualization: Yoánderosn Pérez Díaz, Luis Acao Francois and Yanet Pérez Guerrero.

Writing-original draft: Yoánderosn Pérez Díaz and Rolando Rodríguez Puga.

Writing-review and editing: Yoánderosn Pérez Díaz and Rolando Rodríguez Puga.